



Hematological: Anemia
Reblozyl (luspatercept-aamt) J0896,
Enjaymo (sutimlimab-jome) J1302
Prior Authorization Request
Medicare Part B Form

*Instructions: * Indicates required information – Form may be returned if required information is not provided. Please fax this request to the appropriate fax number listed at the bottom of the page.*

<input type="checkbox"/>	Standard Request– (72 Hours)	<input type="checkbox"/>	Urgent Request (standard time frame could place the member's life, health or ability in serious jeopardy)
Date Requested _____			
Requestor _____ Clinic name: _____ Phone _____ / Fax _____			

MEMBER INFORMATION

*Name: _____ *ID#: _____ *DOB: _____

PRESCRIBER INFORMATION

*Name: _____ MD FNP DO NP PA *Phone: _____

*Address: _____ *Fax: _____

DISPENSING PROVIDER / ADMINISTRATION INFORMATION

*Name: _____ Phone: _____

*Address: _____ Fax: _____

PROCEDURE / PRODUCT INFORMATION

HCPC Code	Name of Drug	Dose (Wt: _____ kg Ht: _____)	Frequency	End Date if known

Self-administered Provider-administered Home Infusion

Chart notes attached. **Other important information:** _____

Diagnosis: ICD10: _____ **Description:** _____

Provider attests the diagnosis provided is an FDA-Approved indication for this drug

CLINICAL INFORMATION

New Start or Initial Request: (Clinical documentation required for all requests)

Reblozyl

Patient has diagnosis of β -thalassemia and ALL of the following:

- Individual is 18 years of age or older; AND
- Individual has a diagnosis of beta thalassemia or hemoglobin E beta (E/ β)-thalassemia; AND
- Documentation is provided that individual required regular red blood cell transfusions at initiation, defined as both of the following (NCT02604433):
 - Individual received six to twenty (6-20) RBC units in the last 24 weeks; AND
 - Individual had no transfusion-free period greater than 35 days in the last 24 weeks; AND
- Individual has a baseline hemoglobin (Hgb) level 11 g/dL or less.

Patient has diagnosis of MDS-RS or MDS/MPN-RS-T or ESA-naïve MDS and ALL of the following:

- Individual is 18 years of age or older; AND
- Individual has one of the following (A, B, or C):

- A. Documentation is provided that individual has a diagnosis very low to intermediate risk myelodysplastic syndromes with ring sideroblasts (MDS-RS) greater than or equal to 15% (or ring sideroblasts 5% to 14% with an SF3B1 mutation) (Label, NCCN 2A); AND
 - Individual meets ONE of the following criteria:
 - Serum erythropoietin (EPO) level of greater than 500 mU/mL; OR
 - Serum EPO level of less than or equal to 500 mU/mL following no response to combination treatment with erythropoiesis-stimulating agent (ESA) and granulocyte-colony stimulating factor (G-CSF); OR
 - B. Individual has a diagnosis of myelodysplastic/myeloproliferative neoplasm with ring sideroblasts and thrombocytosis (MDS/MPN-RS-T) with all of the following:
 - Ring sideroblasts greater than or equal to 15% (WHO 2017), and documentation is provided;
 - Thrombocytosis (defined as platelets greater than or equal to 450 x10⁹/L) (WHO 2017); OR
 - C. Individual has a diagnosis of MDS; AND
 - Documentation is provided that individual has serum EPO level less than 500 U/L;
- Documentation is provided that individual has required regular red blood cell transfusions of two (2) or more RBC units over eight (8) weeks in the last 16 weeks; AND
- Individual has a baseline hemoglobin (Hgb) level 11 g/dL or less.

Enjaymo

- Patient has a diagnosis of cold agglutinin disease (CAD) defined as ALL of the following:
 - The presence of chronic hemolysis;
 - A positive polyspecific direct antiglobulin test result;
 - A monospecific direct antiglobulin test result strongly positive for C3d;
 - A cold agglutinin titer of 1:64 or higher measured at 4oC;
 - A direct antiglobulin test result for IgG of 1+ or less;
 - Presence of one or more symptoms associated with CAD (i.e. symptomatic anemia, acrocyanosis, Raynaud’s phenomenon, hemoglobinuria, disabling circulatory symptoms, or a major adverse vascular event); AND
- Patient is using Enjaymo to decrease the need for red blood cell transfusions due to hemolysis with cold agglutinin disease (CAD)

If not, please provide **clinical rationale** for formulary exception: _____

- Continuation Requests: (Clinical documentation required for all requests)
 - Provider has reviewed the attached “Criteria for Continuation” and attests the member meets ALL required PA Continuation criteria.**
 - Patient had an adequate response or significant improvement while on this medication.
- If not, please provide clinical rationale for continuing this medication: _____

ACKNOWLEDGEMENT

Request By (Signature Required): _____ **Date:** ____ / ____ / ____

Any person who knowingly files a request for authorization of coverage of a medical procedure or service with the intent to injure, defraud or deceive any insurance company by providing materially false information or conceals material information for the purpose of misleading, commits a fraudulent insurance act, which is a crime and subjects such person to criminal and civil penalties. **THIS AUTHORIZATION IS NOT A GUARANTEE OF PAYMENT.** PAYMENT IS BASED ON BENEFITS IN EFFECT AT THE TIME OF SERVICE, MEMBER ELIGIBILITY AND MEDICAL NECESSITY.

Prior Authorization Group – Hematological: Anemia PA

Drug Name(s):

REBLOZYL
ENJAYMO

LUSPATERCEPT-AAMT
SUTIMLIMAB-JOME

Criteria for approval of Prior Authorization Drug:

1. Prescribed for an approved FDA diagnosis (as listed below):
2. Member does not have any clinically relevant contraindications, or CMS/Plan exclusions, to the requested drug.
 - If the member meets all these criteria, they may be approved by the Plan for the requested drug.
 - Quantity limits and Tiering will be determined by the Plan.

Exclusion Criteria:

N/A

Prescriber Restrictions:

N/A

Coverage Duration:

Approvals will be for 6 months

FDA Indications:

Reblozyl

- Anemia, After erythropoiesis stimulating agent failure, requiring 2 or more RBC units over 8 weeks - Myelodysplastic/myeloproliferative neoplasm with ring sideroblasts and thrombocytosis
- Anemia, After erythropoiesis stimulating agent failure, requiring 2 or more RBC units over 8 weeks - Myelodysplastic syndrome, Very low- to intermediate-risk disease with ring sideroblasts (MDS-RS)
- Anemia - Beta thalassemia

Enjaymo

- Cold autoimmune hemolytic anemia - Hemolysis

Off-Label Uses:

N/A

Age Restrictions:

Reblozyl, Enjaymo:

Safety and effectiveness of luspatercept-aamt have not been established in pediatric patients

Other Clinical Considerations:

N/A

Resources:

https://www.micromedexsolutions.com/micromedex2/librarian/CS/2A458B/ND_PR/evidencexpert/ND_P/evidencexpert/DUPLICATI/ONSHIELDSYNC/1D1D55/ND_PG/evidencexpert/ND_B/evidencexpert/ND_AppProduct/evidencexpert/ND_T/evidencexpert/PFActio/nd/evidencexpert.DoIntegratedSearch?SearchTerm=luspatercept&UserSearchTerm=luspatercept&SearchFilter=filterNone&navitem=searchGlobal#



Part B Prior Authorization Guidelines

https://www.micromedexsolutions.com/micromedex2/librarian/CS/077199/ND_PR/evidencexpert/ND_P/evidencexpert/DUPLICATIONSHIELDSYN/C/EEED60/ND_PG/evidencexpert/ND_B/evidencexpert/ND_AppProduct/evidencexpert/ND_T/evidencexpert/PFActionId/evidencexpert.GoToDashboard?docId=933527&contentSetId=100&title=Sutimlimab-jome&servicesTitle=Sutimlimab-jome&brandName=Enjaymo&UserMdxSearchTerm=enjaymo&=null#

CLINICAL / CMS
ONLY